ELSEVIER

Neuroscience Letters 299 (2001) 244–247

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 299

Aerts, L., see Harder, T., 85
Aguilar, J., see Giménez, R., 209
Altememi, G.F., see Mathis, J.P., 173
Anadón, R., Becerra, M., Díaz, M.L., Manso, M.J., Presence and development of thyrotropin-releasing hormone-immunoreactive amacrine cells in the retina of a teleost, the brown trout (*Salmo trutta fario*), 225
Andersen, P.M., see Russ, C., 156
Aoki, T., see Ise, Y., 29
Apkarian, A.V., Krauss, B.R., Fredrickson, B.E., Szeverenyi, N.M., Imaging the pain of low back pain: functional magnetic resonance imaging in combination with monitoring subjective pain perception allows the study of clinical pain states, 57
Arai, N., Furukawa, N., Miyamae, T., Goshima, Y., Sasaki, Y., Ohshima, E., Suzuki, F., Fujita, K., Misu, Y., DOPA cyclohexyl ester, a competitive DOPA antagonist, protects glutamate release and resultant delayed neuron death by transient ischemia in hippocampus CA1 of conscious rats, 213
Asaga, H., Ishigami, A., Protein deimination in the rat brain after kainate administration: citrulline-containing proteins as a novel marker of neurodegeneration, 5
Aust, G., see Reibiger, I., 49

Baker, L.P., see Kumar, P.A., 181
Baker, M., see Russ, C., 156
Banik, R.K., Sato, J., Yajima, H., Mizumura, K., Differences between the Lewis and Sprague–Dawley rats in chronic inflammation induced norepinephrine sensitivity of cutaneous C-fiber nociceptors, 21
Barlovatz-Meimon, G., see Planchenault, T., 140
Bäurle, J., see Radnikow, G., 81
Beart, P.M., see Moldrich, R.X., 125
Becerra, M., see Anadón, R., 225
Bermudez, M.A., see Gonzalez, F., 41
Beyer, R., see Reibiger, I., 49
Bianchi, L., Miller III, D.M., George Jr., A.L., Expression of a ClC chloride channel in *Caenorhabditis elegans* gamma-aminobutyric acid-ergic neurons, 177
Blesa, R., see Queralt, R., 239
Boeck, C.R., see Porciúncula, L.O., 217
Bowden, D.M., see Kumar, P.A., 181
Bullough, J.D., see Rea, M.S., 45

Campbell, K.B., see Muller-Gass, A., 197
Castellví, M., see Queralt, R., 239
Cellini, E., see Nacmias, B., 9
Chapman, A.G., see Moldrich, R.X., 125
Charrière-Bertrand, C., see Planchenault, T., 140
Chen, C.-J., see Liao, S.-L., 113
Cho, H., see Shin, J.S., 135
Cho, S.Y., see Shin, J.S., 135
Coleman, J.R., see Ross, K.C., 234
Condés-Lara, M., Talavera-Cuevas, E., Larriva-Sahd, J., Martínez-Lorenzana, G., Different wheat germ agglutinin-horseradish peroxidase labeling in structures related to the development of amygdaline kindling in the rat, 13
Costa, S., see Planchenault, T., 140
Cruwys, S.C., see Hood, V.C., 201

D'Andrea, F., see Nacmias, B., 9
Díaz-Cabiale, Z., see Razani, H., 145
Diaz, F.G., see Guyot, L.L., 37
Díaz, M.L., see Anadón, R., 225
Dippold, C.S., see Soares, J.C., 150
Dishman, R.K., see O'Neal, H.A., 69
Dudel, J., see Schramm, M., 193

Estacio, M.A.C., see Reyes, B.A.S., 109
Ezquerra, M., see Queralt, R., 239

Fages, C., see Planchenault, T., 140
Fernandes de Lima, V.M., see Peixoto, N.L.V., 89
Ferrer, I., Puig, B., Goutan, E., Gombau, L., Muñoz-Cánoves, P., Methylazoximethanol acetate-induced cell death in the granule cell layer of the developing mouse cerebellum is associated with caspase-3 activation, but does not depend on the tissue-type plasminogen activator, 77
Figueiro, M.G., see Rea, M.S., 45
Fisher, P.J., see Turic, D., 97
Forleo, P., see Nacmias, B., 9
Forster Wells, K., see Soares, J.C., 150
Frank, E., see Soares, J.C., 150
Franke, K., see Harder, T., 85
Fredrickson, B.E., see Apkarian, A.V., 57
Fujita, K., see Arai, N., 213
Fukuyama, N., see Hirabayashi, H., 159
Furlong, R., see Russ, C., 156
Furukawa, N., see Arai, N., 213
Fuxe, K., see Razani, H., 145

Gebhardt, C., see Reibiger, I., 49
George Jr., A.L., see Bianchi, L., 177
Giménez, R., Aguilar, J., Cytidine (5′) diphosphocholine-induced decrease in cerebral platelet activating factor is due to inactivation of its synthesizing enzyme cholinephosphotransferase in aged rats, 209
Gombau, L., see Ferrer, I., 77
Gonzalez, F., Perez, R., Justo, M.S., Bermudez, M.A., Response latencies to visual stimulation and disparity sensitivity in single cells of the awake *Macaca mulatta* visual cortex, 41
Goshima, Y., see Arai, N., 213
Goutan, E., see Ferrer, I., 77

PII: S0304-3940(01)01564-6

Guarnieri, B.M., see Nacmias, B., 9
Guyot, L.L., Diaz, F.G., O'Regan, M.H., McLeod, S., Park, H., Phillis, J.W., Real-time measurement of glutamate release from the ischemic penumbra of the rat cerebral cortex using a focal middle cerebral artery occlusion model, 37

Hager, K., see Massing, W., 101
Hanke, W., see Peixoto, N.L.V., 89
Harder, T., Aerts, L., Franke, K., Van Bree, R., Van Assche, F.A., Plagemann, A., Pancreatic islet transplantation in diabetic pregnant rats prevents acquired malformation of the ventromedial hypothalamic nucleus in their offspring, 85
Harigaya, Y., see Tomidokoro, Y., 169
Hasegaw, Y., Kasai, T., Kinoshita, H., Yahagi, S., Modulation of a motor evoked response to transcranial magnetic stimulation by the activity level of the first dorsal interosseous muscle in humans when grasping a stationary object with different grip widths, 1
Hashimoto, I., see Kimura, T., 61
Hirabayashi, H., Takizawa, S., Fukuyama, N., Nakazawa, H., Shinohara, Y., *N*-methyl-D-aspartate receptor antagonist reduces nitrotyrosine formation in caudate-putamen in rat focal cerebral ischemia-reperfusion, 159
Hirano, M., see Ise, Y., 29
Ho, M., Segre, M., Individual and combined effects of ethanol and cocaine on the human dopamine transporter in neuronal cell lines, 229
Holmes, P.V., see O'Neal, H.A., 69
Hood, V.C., Cruwys, S.C., Urban, L., Kidd, B.L., The neurogenic contribution to synovial leucocyte infiltration and other outcome measures in a guinea pig model of arthritis, 201
Huang, X., Tang, J.-S., Yuan, B., Jia, H., Morphine applied to the ventrolateral orbital cortex produces a naloxone-reversible antinociception in the rat, 189
Hwang, S.W., see Shin, J.S., 135

I'Anson, H., see Reyes, B.A.S., 109
Ichikawa, T., Mutual coupling among insect neurosecretory cells with an ultradian firing rhythm, 73
Ikeda, M., see Tomidokoro, Y., 169
Ise, Y., Katayama, S., Hirano, M., Aoki, T., Narita, M., Suzuki, T., Effects of fluvoxamine on morphine-induced inhibition of gastrointestinal transit, antinociception and hyperlocomotion in mice, 29
Ishigami, A., see Asaga, H., 5
Ishiguro, K., see Tomidokoro, Y., 169
Ishizaki, M., Oka, Y., Amperometric recording of gonadotropin-releasing hormone release activity in the pituitary of the dwarf gourami (teleosat) brain-pituitary slices, 121
Isobe, K., Kusaka, T., Nagano, K., Okubo, K., Yasuda, S., Kondo, M., Itoh, S., Onishi, S., Functional imaging of the brain in sedated newborn infants using near infrared topography during passive knee movement, 221
Itoh, S., see Isobe, K., 221

Jellinger, K.A., see Rösler, N., 117
Jia, H., see Huang, X., 189
Joe, E.-h., see Kang, J., 205
Jones, L.S., see Ross, K.C., 234
Jou, I., see Kang, J., 205
Justo, M.S., see Gonzalez, F., 41

Kaba, H., see Osako, Y., 65
Kaneko, T., see Li, Y.-Q., 33
Kang, J., Yang, M.-s., Jou, I., Joe, E.-h., Identification of protein kinase C isoforms involved in interferon-gamma-induced expression of inducible nitric oxide synthase in murine BV2 microglia, 205
Kanno, I., see Kashikura, K., 53
Kasai, T., see Hasegaw, Y., 1
Kashikura, A., see Kashikura, K., 53
Kashikura, K., Kershaw, J., Kashikura, A., Zhang, X., Matsuura, T., Kanno, I., Hyperoxia modified activation-induced blood oxygenation level-dependent response of human visual cortex (V1): an event-related functional magnetic resonance imaging study, 53
Katayama, S., see Ise, Y., 29
Kawarabayashi, T., see Tomidokoro, Y., 169
Kenklies, M., see Massing, W., 101
Kershaw, J., see Kashikura, K., 53
Kidd, B.L., see Hood, V.C., 201
Kimura, T., Hashimoto, I., Source of somatosensory primary cortical evoked magnetic fields (N20m) elicited by index finger stimulation moves toward mediolateral direction in area 3b in man, 61
Kinoshita, H., see Hasegaw, Y., 1
Kondo, M., see Isobe, K., 221
Krauss, B.R., see Apkarian, A.V., 57
Kumar, P.A., Baker, L.P., Storm, D.R., Bowden, D.M., Expression of type I adenylyl cyclase in intrinsic pathways of the hippocampal formation of the macaque (*Macaca nemestrina*), 181
Kupfer, D.J., see Soares, J.C., 150
Kusaka, T., see Isobe, K., 221
Kwon, M.J., see Shin, J.S., 135

Larriva-Sahd, J., see Condés-Lara, M., 13
Lee, S.-Y., see Shin, J.S., 135
Li, Y.-Q., Kaneko, T., Mizuno, N., Collateral projections of nucleus raphe dorsalis neurones to the caudate-putamen and region around the nucleus raphe magnus and nucleus reticularis gigantocellularis pars α in the rat, 33
Liao, S.-L., Chen, C.-J., Differential effects of cytokines and redox potential on glutamate uptake in rat cortical glial cultures, 113
Lleó, A., see Queralt, R., 239
Logan, J., see Muller-Gass, A., 197
Löscher, W., see Redecker, P., 93
Lovestone, S., see Russ, C., 156
Ludolph, A.C., see von Arnim, C.A.F., 130

Ma, N., see Yasuda, E., 162
Mades, S., see Radnikow, G., 81
Maeda, K.-i., see Reyes, B.A.S., 109
Mallinger, A.G., see Soares, J.C., 150
Mandyam, C.D., see Mathis, J.P., 173
Mann, D., see Russ, C., 156
Manso, M.J., see Anadón, R., 225
Marcoux, A., see Muller-Gass, A., 197
Martínez-Lorenzana, G., see Condés-Lara, M., 13
Massing, W., Kenklies, M., Hager, K., Virtual image grasping in patients with Alzheimer-type dementia, 101
Mathis, J.P., Mandyam, C.D., Altememi, G.F., Pasternak, G.W., Standifer, K.M., Orphanin FQ/nociceptin and naloxone benzoylhydrazone activate distinct receptors in BE(2)-C human neuroblastoma cells, 173
Matsubara, E., see Tomidokoro, Y., 169
Matsuura, T., see Kashikura, K., 53
McLeod, S., see Guyot, L.L., 37

Mehta, H., Thomas, B., Mohanakumar, K.P., Effects of *p*-chlorophenylalanine on striatal acetylcholinesterase activity and on biogenic amine levels in nuclei raphe and caudate-putamen during physostigmine-induced tremor in rats, 105
Meldrum, B.S., see Moldrich, R.X., 125
Miller III, D.M., see Bianchi, L., 177
Misane, I., see Razani, H., 145
Misgeld, U., see Radnikow, G., 81
Misu, Y., see Arai, N., 213
Miyamae, T., see Arai, N., 213
Mizumura, K., see Banik, R.K., 21
Mizumura, K., see Sato, J., 17
Mizuno, N., see Li, Y.-Q., 33
Mohanakumar, K.P., see Mehta, H., 105
Moldrich, R.X., Talebi, A., Beart, P.M., Chapman, A.G., Meldrum, B.S., The $mGlu_{2/3}$ agonist 2*R*,4*R*-4-aminopyrrolidine-2,4-dicarboxylate, is anti- and proconvulsant in DBA/2 mice, 125
Muller-Gass, A., Marcoux, A., Logan, J., Campbell, K.B., The intensity of masking noise affects the mismatch negativity to speech sounds in human subjects, 197
Muñoz-Cánoves, P., see Ferrer, I., 77

Nacmias, B., Tedde, A., Cellini, E., Forleo, P., Orlacchio, A., Guarnieri, B.M., Petruzzi, C., D'Andrea, F., Serio, A., Sorbi, S., α2-Macroglobulin polymorphisms in Italian sporadic and familial Alzheimer's disease, 9
Nagano, K., see Isobe, K., 221
Nakazawa, H., see Hirabayashi, H., 159
Narita, M., see Ise, Y., 29

Ögren, S.O., see Razani, H., 145
Oh, U., see Shin, J.S., 135
Ohshima, E., see Arai, N., 213
Oka, T., see Osako, Y., 65
Oka, Y., see Ishizaki, M., 121
Okamoto, K., see Tomidokoro, Y., 169
Okubo, K., see Isobe, K., 221
Oliva, R., see Queralt, R., 239
Omura, S., see Sato, J., 17
O'Neal, H.A., Van Hoomissen, J.D., Holmes, P.V., Dishman, R.K., Prepro-galanin messenger RNA levels are increased in rat locus coeruleus after treadmill exercise training, 69
Onishi, S., see Isobe, K., 221
O'Regan, M.H., see Guyot, L.L., 37
Orlacchio, A., see Nacmias, B., 9
Osako, Y., Otsuka, T., Taniguchi, M., Oka, T., Kaba, H., Oxytocin enhances presynaptic and postsynaptic glutamatergic transmission between rat olfactory bulb neurones in culture, 65
Otsuka, T., see Osako, Y., 65
Owen, M.J., see Turic, D., 97

Pabst, H., see Redecker, P., 93
Park, H., see Guyot, L.L., 37
Park, J.-M., see Tomidokoro, Y., 169
Pasternak, G.W., see Mathis, J.P., 173
Peixoto, N.L.V., Fernandes de Lima, V.M., Hanke, W., Correlation of the electrical and intrinsic optical signals in the chicken spreading depression phenomenon, 89
Perez, R., see Gonzalez, F., 41
Perzelova, A., see Planchenault, T., 140
Petruzzi, C., see Nacmias, B., 9
Phillis, J.W., see Guyot, L.L., 37
Pickering-Brown, S.M., see Russ, C., 156
Plagemann, A., see Harder, T., 85
Planchenault, T., Costa, S., Fages, C., Riche, D., Charrière-Bertrand, C., Perzelova, A., Barlovatz-Meimon, G., Tardy, M., Differential expression of laminin and fibronectin and of their related metalloproteinases in human glioma cell lines: relation to invasion, 140
Plomin, R., see Turic, D., 97
Porciúncula, L.O., Rocha, J.B.T., Boeck, C.R., Vendite, D., Souza, D.O., Ebselen prevents excitotoxicity provoked by glutamate in rat cerebellar granule neurons, 217
Powell, J.F., see Russ, C., 156
Puig, B., see Ferrer, I., 77

Queralt, R., Ezquerra, M., Castellví, M., Lleó, A., Blesa, R., Oliva, R., Detection of the *presenilin 1* gene mutation (M139T) in early-onset familial Alzheimer disease in Spain, 239

Radnikow, G., Titz, S., Mades, S., Bäurle, J., Misgeld, U., γ-Aminobutyric $acid_B$ autoreceptors in substantia nigra and neostriatum of the weaver mutant mouse, 81
Razani, H., Díaz-Cabiale, Z., Misane, I., Wang, F.-H., Fuxe, K., Ögren, S.O., Prolonged effects of intraventricular galanin on a 5-hydroxytryptamine$_{1A}$ receptor mediated function in the rat, 145
Rea, M.S., Bullough, J.D., Figueiro, M.G., Human melatonin suppression by light: a case for scotopic efficiency, 45
Redecker, P., Pabst, H., Löscher, W., Steinlechner, S., Evidence for microvesicular storage and release of glycine in rodent pinealocytes, 93
Reibiger, I., Aust, G., Tscheudschilsuren, G., Beyer, R., Gebhardt, C., Spanel-Borowski, K., The expression of substance P and its neurokinin-1 receptor mRNA in the bovine corpus luteum of early developmental stage, 49
Reyes, B.A.S., Estacio, M.A.C., I'Anson, H., Tsukamura, H., Maeda, K.-i., Glucoprivation increases estrogen receptor α immunoreactivity in the brain catecholaminergic neurons in ovariectomized rats, 109
Riche, D., see Planchenault, T., 140
Riepe, M.W., see von Arnim, C.A.F., 130
Rocha, J.B.T., see Porciúncula, L.O., 217
Rösler, N., Wichart, I., Jellinger, K.A., Ex vivo lumbar and post mortem ventricular cerebrospinal fluid substance P-immunoreactivity in Alzheimer's disease patients, 117
Ross, K.C., Coleman, J.R., Jones, L.S., Anti-epileptiform effects of audiogenic seizure priming on in vitro kindling in rat hippocampus, 234
Russ, C., Lovestone, S., Baker, M., Pickering-Brown, S.M., Andersen, P.M., Furlong, R., Mann, D., Powell, J.F., The extended haplotype of the microtubule associated protein tau gene is not associated with Pick's disease, 156

Sasaki, Y., see Arai, N., 213
Sato, J., see Banik, R.K., 21
Sato, J., Takanari, K., Omura, S., Mizumura, K., Effects of lowering barometric pressure on guarding behavior, heart rate and blood pressure in a rat model of neuropathic pain, 17
Schramm, M., Dudel, J., Pertussis toxin does not affect the time course of quantal release in crayfish and mouse muscle, but has other post- and presynaptic effects, especially on adenosine autoreceptors, 193
Schrauf, M., Stern, C., The visual resolution of Landolt-C optotypes in human subjects depends on their orientation: the 'gap-down' effect, 185
Segre, M., see Ho, M., 229
Semba, R., see Yasuda, E., 162

Serio, A., see Nacmias, B., 9
Shin, J.S., Wang, M.-H., Hwang, S.W., Cho, H., Cho, S.Y., Kwon, M.J., Lee, S.-Y., Oh, U., Differences in sensitivity of vanilloid receptor 1 transfected to human embryonic kidney cells and capsaicin-activated channels in cultured rat dorsal root ganglion neurons to capsaicin receptor agonists, 135
Shinohara, Y., see Hirabayashi, H., 159
Shoji, M., see Tomidokoro, Y., 169
Soares, J.C., Dippold, C.S., Forster Wells, K., Frank, E., Kupfer, D.J., Mallinger, A.G., Increased platelet membrane phosphatidylinositol-4,5-bisphosphate in drug-free depressed bipolar patients, 150
Sorbi, S., see Nacmias, B., 9
Souza, D.O., see Porciúncula, L.O., 217
Spanel-Borowski, K., see Reibiger, I., 49
Standifer, K.M., see Mathis, J.P., 173
Steinlechner, S., see Redecker, P., 93
Stern, C., see Schrauf, M., 185
Storm, D.R., see Kumar, P.A., 181
Suzuki, F., see Arai, N., 213
Suzuki, T., see Ise, Y., 29
Szeverenyi, N.M., see Apkarian, A.V., 57

Takanari, K., see Sato, J., 17
Takizawa, S., see Hirabayashi, H., 159
Talavera-Cuevas, E., see Condés-Lara, M., 13
Talebi, A., see Moldrich, R.X., 125
Tang, J.-S., see Huang, X., 189
Taniguchi, M., see Osako, Y., 65
Tardy, M., see Planchenault, T., 140
Tedde, A., see Nacmias, B., 9
Thomas, B., see Mehta, H., 105
Timmler, M., see von Arnim, C.A.F., 130
Titz, S., see Radnikow, G., 81
Tomidokoro, Y., Ishiguro, K., Harigaya, Y., Matsubara, E., Ikeda, M., Park, J.-M., Yasutake, K., Kawarabayashi, T., Okamoto, K., Shoji, M., Aβ amyloidosis induces the initial stage of tau accumulation in APP_{Sw} mice, 169
Tscheudschilsuren, G., see Reibiger, I., 49
Tso, P.H., Wong, Y.H., Opioid-induced adenylyl cyclase supersensitization in human embryonic kidney 293 cells requires pertussis toxin-sensitive G proteins other than G_{i1} and G_{i3}, 25
Tsukamura, H., see Reyes, B.A.S., 109
Turic, D., Fisher, P.J., Plomin, R., Owen, M.J., No association between apolipoprotein E polymorphisms and general cognitive ability in children, 97

Ulfig, N., Expression of leukemia inhibitory factor in the ganglionic eminence of the human fetal brain after bleedings, 153
Urban, L., see Hood, V.C., 201

Van Assche, F.A., see Harder, T., 85
Van Bree, R., see Harder, T., 85
Van Hoomissen, J.D., see O'Neal, H.A., 69
Vendite, D., see Porciúncula, L.O., 217
von Arnim, C.A.F., Timmler, M., Ludolph, A.C., Riepe, M.W., Chemical preconditioning in mice is not mediated by upregulation of nitric oxide synthase isoforms, 130

Wang, F.-H., see Razani, H., 145
Wang, M.-H., see Shin, J.S., 135
Wichart, I., see Rösler, N., 117
Wong, Y.H., see Tso, P.H., 25

Yahagi, S., see Hasegaw, Y., 1
Yajima, H., see Banik, R.K., 21
Yang, M.-s., see Kang, J., 205
Yasuda, E., Ma, N., Semba, R., Immunohistochemical evidences for localization and production of D-serine in some neurons in the rat brain, 162
Yasuda, S., see Isobe, K., 221
Yasutake, K., see Tomidokoro, Y., 169
Yuan, B., see Huang, X., 189

Zhang, X., see Kashikura, K., 53